



## I. Epidemiological profile

Population (UN Population Division)	2017	%	Parasites and vectors
Number of active foci	18	-	<b>Major plasmodium species:</b> <b>P.falciparum: 100 (%) , P.vivax: 0 (%)</b>
Number of people living within active foci	175.7K	32	Major anopheles species: <i>An. arabiensis</i>
Malaria free (0 cases)	370.7K	68	
Total	370.7K		
			Reported indigenous confirmed cases (health facility): 423
			Confirmed cases at community level: -
			Confirmed cases from private sector: -
			Indigenous deaths: 1

## II. Intervention policies and strategies

Intervention	Policies/Strategies	Yes/No	Year adopted	Antimalaria treatment policy	Medicine	Year adopted						
ITN	ITNs/LLINs distributed free of charge	No	-	First-line treatment of unconfirmed malaria	AL	2007						
	ITNs/LLINs distributed to all age groups	No	-	First-line treatment of <i>P. falciparum</i>	AL	2007						
IRS	IRS is recommended	Yes	1998	For treatment failure of <i>P. falciparum</i>	QN	-						
	DDT is used for IRS	No	-	Treatment of severe malaria	QN	-						
Larval control	Use of Larval Control	Yes	-	Treatment of <i>P. vivax</i>	-	-						
IPT	IPT used to prevent malaria during pregnancy	No	-	Dosage of primaquine for radical treatment of <i>P. vivax</i>								
				Type of RDT used	Pf + all species (Combo)							
Diagnosis	Patients of all ages should receive diagnostic test	Yes	1998	Therapeutic efficacy tests (clinical and parasitological failure, %)								
	Malaria diagnosis is free of charge in the public sector	Yes	1975	Medicine	Year	Min	Median	Max	Follow-up	No. of studies	Species	
Treatment	ACT is free for all ages in public sector	Yes	2008	Resistance status by insecticide class (2010-2017) and use of class for malaria vector control (2017)								
	The sale of oral artemisinin-based monotherapies (oAMTs) are allowed		-	Insecticide class	Years	(%) sites <sup>1</sup>	Vectors <sup>2</sup>	Used <sup>3</sup>				
	Single dose of primaquine (0.25 mg base/kg) is used as gametocidal medicine for <i>P. falciparum</i>	Yes	-	Carbamates	2017-2017	0% (1)	-	No				
	Primaquine is used for radical treatment of <i>P. vivax</i>	No	2015	Organochlorines	-	-	-	No				
	G6PD test is a requirement before treatment with primaquine	No	-	Organophosphates	2017-2017	0% (1)	-	No				
	Directly observed treatment with primaquine is undertaken	Yes	-	Pyrethroids	2017-2017	100% (1)	<i>An. gambiae s.l.</i>	Yes				
	System for monitoring of adverse reaction to antimalarials exists	No	2007	<sup>1</sup> Percent of sites for which resistance confirmed and total number of sites that reported data (n)								
	Surveillance	ACD for case investigation (reactive)	Yes	2001	<sup>2</sup> Principal vectors that exhibited resistance							
	ACD at community level of febrile cases (pro-active)	Yes	2001	<sup>3</sup> Class used for malaria vector control in 2017								
	Mass screening is undertaken	No	2001									
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	Yes	2007									
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	-									
	Case and foci investigation undertaken	Yes	-									
	Case reporting from private sector is mandatory	Yes	-									